



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/14	A1	(11) International Publication Number: WO 96/38131
		(43) International Publication Date: 5 December 1996 (05.12.96)

(21) International Application Number: PCT/EP96/02299

(22) International Filing Date: 30 May 1996 (30.05.96)

(30) Priority Data:
9511220.7 2 June 1995 (02.06.95) GB

(71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): BUTLER, James, Matthew [GB/GB]; Glaxo Research and Development Limited, Park Road, Ware, Hertfordshire SG12 0DP (GB).

(74) Agents: GALLAFENT, Alison et al.; Glaxo Wellcome plc, Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: METHOD OF PRODUCING A SOLID DISPERSION OF A POORLY WATER SOLUBLE DRUG

(57) Abstract

Solid dispersions of poorly soluble drugs, to processes for their preparation and their use in pharmaceutical compositions. Specifically, solid dispersions of (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione and (+)-N-[1-(adamantanmethyl)-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro-1H-1,5benzodiazepin-3-yl]-N'-phenylurea are described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore -
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

METHOD OF PRODUCING A SOLID DISPERSION OF A POORLY WATER SOLUBLE DRUG

The present invention relates to the field of solid dispersions of poorly water soluble drugs, to processes for their preparation and their use in pharmaceutical compositions. More particularly the present invention relates to solid dispersions in the form of co-precipitates of poorly water soluble drugs and their compositions with a pharmaceutically acceptable carrier or excipient therefor. Specifically, the invention relates to co-precipitates of (a) a potent and selective inhibitor of cyclic guanosine 3',5'-monophosphate specific phosphodiesterase (cGMP specific PDE) and (b) a potent and selective gastrin and CCK B antagonist, processes for the preparation of such solid dispersions, pharmaceutical compositions containing the same and their use thereof in therapy.

Co-precipitation is a recognised technique for increasing the dissolution of poorly water soluble drugs, such as griseofulvin, ketoprofen, sulphathiazide, spirinolactone, tolbutamide and nifedipine, so as to consequently improve bioavailability thereof. Techniques such as solvent deposition, lyophilization, solvate formation and solid dispersion (of which co-precipitation is an example as described above) have therefore been developed to try to overcome the problem of poor water solubility and resultant low bioavailability.

Solid dispersions in the pharmaceutical field are dispersions of one or more active ingredients, generally poorly water soluble drugs, in an inert carrier or matrix at solid state, which are prepared by either melting the two (fusion), or dissolving them in a solvent, or a combination of approaches, followed by removal of the solvent.

Manufacture of pharmaceutical dispersions by the above referred to melting or fusion technique, involves fusion of the two components where the drug and the carrier are allowed to melt at temperatures at or above the melting point of both the drug and carrier. In the fusion process, the drug and carrier are first blended and both melted in a suitable mixer. The molten mixture is then cooled rapidly to provide a congealed mass which is subsequently milled to produce a powder. The fusion process is technically simple provided that the drug and

carrier are miscible in the molten state but this is not always the case and furthermore, the process is limited in that it tends to lead to drug decomposition due to the high temperatures required to melt the two components.

5 The solvent-based process uses organic solvents to dissolve and intimately disperse the drug and carrier molecules. Identification of a common solvent for both drug and carrier can be problematic, and complete solvent removal from the product can be a lengthy process. In addition, large volumes of solvents are generally required which can give rise to toxicological problems. The drug and
10 carrier are typically dissolved in a solvent such as methylene chloride, acetone, ethanol and mixtures thereof and the solvent is later removed by precipitation techniques, evaporation or the like, while the drug/carrier solid dispersion is collected as a powdered mass.

15 In the case where there is difficulty with thermal instability and immiscibility between the drug and the carrier, the hybrid fusion-solvent method can be employed. The drug is first dissolved in a small quantity of organic solvent and added to the molten carrier. The solvent is then evaporated to generate a product that is subsequently milled to produce a powder. The
20 pharmacokinetics, dissolution rates and processes for formulation of many different solid pharmaceutical dispersions is discussed at length in an article by Ford J., in *Pharm. Acta. Helv.* 61, 3; 69 - 88 (1986).

25 Co-precipitation techniques employ the use of an organic solvent or solvents to dissolve and intimately disperse the drug and carrier molecules as hereinbefore described. Separation of the drug and carrier from the solvent on precipitation can rely on the solubility properties of either the drug or carrier. For example, Simonelli et al, *Journal of Pharmaceutical Sciences*, Vol. 58, No. 5, May 1969, describes a co-precipitation process wherein sulfathiazole is dissolved in
30 sodium hydroxide, followed by addition of polyvinylpyrrolidone; hydrochloric acid is then added to effect co-precipitation. This process is based on co-precipitation employing the solubility of the drug at different pH values. Such reliance on the solubility of the drug may be problematic in that it is not generally applicable to poorly water soluble drugs, as many such drugs do not
35 exhibit a pH dependent solubility. Florence et al, *Communications, J. Pharm.*

Pharmac., 1976, 28 601, describes co-precipitation of trifluoperazine embonate and the polymers poly DL-aspartic acid and polymethylmethacrylate. The co-precipitates were prepared by dissolving the drug and polymer in dimethylformamide and adding the solution to a rapidly stirred volume of water.
5 Both polymers and drug are insoluble in water.

In general terms, problems which can be associated with known co-precipitation techniques can include excess solvent usage, identifying carrier/drug combinations which can be effectively precipitated and enhance bioavailability,
10 the use of heat to effect solution which may detrimentally affect the drug, and the like. Co-precipitation techniques are however attractive for the preparation of solid dispersions, in that less solvents and heat are employed when compared to techniques such as co-evaporation and solvent removal may therefore be facilitated.

15 We have now developed a co-precipitation technique which alleviates the above described disadvantages associated with known techniques, and have also found that co-precipitation offers an advantageous preparation route for solid dispersions of poorly water soluble drugs.

20 There is therefore provided in a first aspect of the present invention a process of preparing a solid dispersion comprising a poorly water soluble drug or salts or solvates (e.g. hydrates) thereof, and a pharmaceutically acceptable carrier or excipient therefor, which process comprises:

- 25 (i) providing an intimate mixture comprising the carrier or excipient and a non-aqueous, water miscible solvent or combination of solvents, and optionally, water;
- 30 (ii) co-mixing the intimate mixture obtained in step (a) with a poorly water soluble drug; and
- (iii) co-precipitating the poorly water soluble drug and the carrier or excipient.

35

As used herein, the term "intimate mixture" can denote a solution, suspension, emulsion, colloid, dispersion or the like. Generally, the term "intimate mixture" as used herein denotes a solution.

5 It has been found surprisingly that small amounts of water in the intimate mixture can aid dissolution of the subsequently added poorly water soluble drug. For example, a 10% ratio of water in a solvent may aid dissolution of the poorly water soluble drug.

10 In a further aspect, the invention describes a process of preparing a solid dispersion comprising a particular cGMP specific PDE (PDEV) inhibitor. More particularly, co-precipitation overcomes problems associated with other preparatory processes for formulating the subject PDEV inhibitor.

15 There is therefore provided by the present invention a process of preparing a solid dispersion comprising (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (hereinafter referred to as Compound A) or salts or solvates (e.g. hydrates) thereof, and a pharmaceutically acceptable carrier or excipient therefor, which
20 process comprises co-precipitating Compound A and the pharmaceutically acceptable carrier or excipient.

In a yet further aspect, the invention describes a process of preparing a solid dispersion comprising a particular gastrin and CCK-B antagonist. More
25 particularly, co-precipitation overcomes problems associated with other preparatory processes for formulating the subject gastrin and CCK-B antagonist.

There is therefore provided by the present invention a process of preparing a solid dispersion comprising (+)-N-[1-(Adamantanmethyl)-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro - 1H-1,5benzodiazepin-3-yl]-N'-phenylurea -(hereinafter
30 referred to as Compound B) or salts or solvates (e.g. hydrates) thereof, and a pharmaceutically acceptable carrier or excipient therefor, which process comprises co-precipitating Compound B and the pharmaceutically acceptable carrier or excipient. The synthesis and use of Compound B has been previously
35 described in WO93.14074.

Suitably the co-precipitation of Compound A or B comprises the steps of:

- 5 (a) providing an intimate mixture of a poorly water soluble drug selected from Compound A or Compound B, the carrier or excipient therefor and a non-aqueous, water miscible solvent or combination of solvents, and optionally, water; and
- 10 (b) co-precipitating the compound and the carrier or excipient.

It is generally advantageous in step (a) to first co-mix the carrier or excipient together with the solvent or solvents and optional water, thereby providing an initial intimate mixture, prior to addition of the poorly water soluble drug thereto. Subsequently, the drug can be added to the initial intimate mixture. Optionally,

15 the carrier or excipient and solvent initial intimate mixture can be subjected to heating sufficient to facilitate dissolving of the former in the latter. Such a sequence of steps (also substantially as hereinbefore described according to the first aspect of the present invention) can be beneficial in allowing the employ of heat to effect dissolving, whilst obviating any detrimental affect by the heat on

20 the drug.

A co-precipitation step, substantially as described in step (iii), or substantially as described in step (b), can aptly comprise adding the drug, carrier or excipient and solvent to a co-precipitation medium in which the carrier or excipient is

25 insoluble. The resultant co-precipitate can be separated from the remaining components, suitably by filtering or the like, and the co-precipitate washed to remove residual solvent, and dried. The co-precipitate can then be formulated in a suitable pharmaceutical form employing known formulatory techniques, substantially as hereinafter described.

30 The carrier or excipient for the drug, and the co-precipitation medium, are respectively chosen so that the carrier or excipient is substantially insoluble in the co-precipitation medium. It is also advantageous that the carrier or excipient has selected dissolution properties in vivo; for example, the carrier or excipient

35 may be such so as to dissolve rapidly (within about 15 to 60 minutes) in vivo,

alternatively the carrier or excipient may be such so as to dissolve over a relatively prolonged period of time (typically, 2 to 4 hours) so as to achieve sustained release of drug in vivo.

- 5 Suitable carrier or excipients include pharmaceutically acceptable polymeric materials, typical examples being hydroxypropyl methyl cellulose phthalate, polymethylacrylate, hydroxypropyl cellulose and other like carrier or excipient materials. Particularly preferred is hydroxypropyl methyl cellulose phthalate as a carrier or excipient, and there is further provided by the present invention co-precipitates consisting of Compound A and hydroxypropyl methyl cellulose phthalate, and Compound B and hydroxypropyl methyl cellulose phthalate.
- 10

- 15 Aptly the co-precipitation medium comprises an aqueous medium, which is optimally such that the carrier or excipient is substantially insoluble therein as substantially hereinbefore described.

- 20 Conveniently the following combinations of carrier or excipient and co-precipitation medium can be employed in a process according to the present invention:

- 25 (a) in the case where the carrier or excipient is hydroxypropyl methyl cellulose phthalate, the co-precipitation medium is suitably a weakly acidic medium (pH in the range of 0.5 to 5.0, typically 0.8 to 2.0), typically 0.5N hydrochloric acid or acetic acid;

- 30 (b) in the case where the carrier or excipient is an acid soluble polymethylacrylate, the co-precipitation medium is suitably a neutral or basic medium, (pH in the range 6.0 to 13.0), water or dilute alkali being appropriate representatives of suitable co-precipitation media; and

- 35 (c) in the case where the carrier or excipient is hydroxypropylcellulose, an appropriate co-precipitation medium is again water, aptly with a temperature of greater than about 40°C, such as 70 to 80°C.

It is of course envisaged that other suitable combinations of carrier or excipient and co-precipitation media may be employed, as will be envisaged by a person skilled in the art. A particularly appropriate combination of carrier or excipient and co-precipitation medium, is hydroxypropyl methyl cellulose phthalate and a dilute acidic medium, substantially as hereinbefore described in point (a) above.

Appropriately, the solvent employed in a process according to the present invention is selected from the group consisting of acetone, methanol, dimethylacetamide, dimethylsulphoxide, dimethylformamide, tetrahydrofuran, and combinations thereof, and optionally, water, although other suitable solvents could be employed. Generally 9:1 acetone/water, 9:1 tetrahydrofuran/water, or 1:1 acetone/methanol mixtures are employed in a process according to the present invention.

There is still further provided by the present invention a solid dispersion consisting essentially of Compound A or Compound B and a pharmaceutically acceptable carrier or excipient therefor. As a further aspect of the present invention, there is provided hereby a solid dispersion comprising Compound A or Compound B obtainable by a process substantially as hereinbefore described.

It has been shown that Compound A is a potent and selective inhibitor of PDEV. Thus, Compound A is of interest for use in therapy, specifically for the treatment of a variety of conditions where inhibition of PDEV is thought to be beneficial.

As a consequence of the selective PDEV inhibition exhibited, cGMP levels are elevated, which in turn can give rise to beneficial anti-platelet, anti-neutrophil, anti-vasospastic, vasodilatory, natriuretic and diuretic activities as well as potentiation of the effects of endothelium-derived relaxing factor (EDRF), nitrovasodilators, atrial natriuretic factor (ANF), brain natriuretic peptide (BNP), C-type natriuretic peptide (CNP) and endothelium-dependent relaxing agents such as bradykinin, acetylcholine and 5-HT₁. Elevated cGMP levels may also mediate relaxation of the corpus cavernosum tissue and consequent penile erection in the treatment of male sexual dysfunction. The solid dispersions of

Compound A therefore have utility in the treatment of a number of disorders, including stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, renal failure, atherosclerosis, conditions of reduced blood vessel patency (e.g. post-percutaneous
5 transluminal coronary angioplasty), peripheral vascular disease, vascular disorders such as Raynaud's disease, inflammatory diseases, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis, glaucoma, diseases characterised by disorders of gut motility (e.g. irritable bowel syndrome), and symptoms associated with male sexual dysfunction.

10 It will be appreciated that references herein to treatment extend to prophylaxis as well as treatment of established conditions.

There is thus provided as a further aspect of the invention a solid dispersion of
15 Compound A substantially as hereinbefore described for use in therapy, in particular for use in the treatment of conditions where inhibition of PDEV is of therapeutic benefit.

According to another aspect of the invention, there is provided the use of a solid
20 dispersion of Compound A substantially as hereinbefore described for the manufacture of a medicament for use in therapy, in particular the treatment of conditions where inhibition of PDEV is of therapeutic benefit.

In a further aspect, the invention provides a method of treating conditions where
25 inhibition of PDEV is of therapeutic benefit in a human or non-human animal body which comprises administering to said body a therapeutically effective amount of a solid dispersion of Compound A substantially as hereinbefore described.

For administration to man in the curative or prophylactic treatment of the
30 disorders identified above, oral dosages of Compound A will generally be in the range of from 0.5-800mg daily for an average adult patient (70kg). Thus for a typical adult patient, individual tablets or capsules contain from 0.2-400mg of active compound, in a suitable pharmaceutically acceptable vehicle or carrier,
35 for administration in single or multiple doses, once or several times per day.

Dosages for intravenous, buccal or sublingual administration will typically be within the range of from 0.1-400 mg per single dose as required. In practice the physician will determine the actual dosing regimen which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case but there can be individual instances in which higher or lower dosage ranges may be merited, and such are within the scope of this invention.

It has been shown that Compound B is a potent and specific antagonist of gastrin and/or CCK-B. The compound has been shown to be an antagonist of CCK, particularly at CCK-B receptors as demonstrated for example by the compound's ability to inhibit the contractile actions of CCK-4 in the presence of a CCK-A antagonist, in the guinea-pig isolated ileum longitudinal muscle-myenteric plexus.

A solid dispersion of Compound B is therefore useful for the treatment and/or prevention of disorders in mammals, especially humans, where modification of the effects of gastrin or CCK-B is of therapeutic benefit. Thus the solid dispersion of Compound B is useful for the treatment of central nervous system disorders where CCK-B and/or gastrin are involved, for example anxiety disorders (including panic disorder, agoraphobia, social phobia, simple phobia, obsessive compulsive disorders, post traumatic stress disorder, and general anxiety disorder), tardive dyskinesia, depression, Parkinson's disease or psychosis. The solid dispersion of Compound B is also useful for the treatment of gastrointestinal disorders especially those where there is an advantage in lowering gastric acidity. Such disorders include peptic ulceration, reflux oesophagitis and Zollinger Ellison syndrome. It may also be useful for the treatment of gastrointestinal disorders such as irritable bowel syndrome, excess pancreatic secretion, acute pancreatitis, motility disorders, antral G cell hyperplasia, fundic mucosal hyperplasia or gastrointestinal neoplasms. It may also be useful for the treatment of dependency on drugs or substances of abuse and withdrawal, Gilles de la Tourette syndrome, or dysfunction of appetite regulatory systems; as well as the treatment of certain tumours of the lower oesophagus, stomach, intestines and colon. Compounds of the invention are

also useful for directly inducing analgesia, or enhancing opiate or non-opiate mediated analgesia, as well as anaesthesia or loss of the sensation of pain.

5 There is thus provided as a further aspect of the invention a solid dispersion of compound B substantially as hereinbefore defined for use in the treatment of conditions where modification of the effects of gastrin and/or CCK-B is of therapeutic benefit.

10 According to another aspect the invention provides the use of a solid dispersion of compound B substantially as hereinbefore defined for the manufacture of a medicament for use in therapy, in particular for the treatment of conditions where modification of the effects of gastrin and/or CCK-B is of therapeutic benefit.

15 According to a further aspect of the invention we provide a method for the treatment of a mammal, including man, in particular in the treatment of conditions where modification of the effects of gastrin and/or CCK-B is of therapeutic benefit which method comprises administering an effective amount of a solid dispersion of Compound B substantially as hereinbefore defined to the
20 patient.

It will be appreciated that the amount of Compound B required for use in treatment will vary with the nature of the condition being treated and the age and the condition of the patient and will be ultimately at the discretion of the
25 attendant physician or veterinarian. In general however doses employed for adult human treatment will typically be in the range of 0.01-2000mg per day e.g 0.01-500mg per day.

30 For human use, a solid dispersion substantially as hereinbefore described will generally be administered in admixture with a pharmaceutical carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, the compound may be administered orally in the form of tablets containing excipients such as cellulose or lactose, or in capsules or
35 ovules either alone or in admixture with excipients.

Thus, the invention provides in a further aspect a pharmaceutical composition comprising a solid dispersion substantially as hereinbefore described together with a pharmaceutically acceptable carrier therefor.

5 There is further provided by the present invention a process of preparing a pharmaceutical composition comprising a solid dispersion substantially as hereinbefore described, which process comprises mixing a solid dispersion substantially as hereinbefore described together with a pharmaceutically acceptable carrier therefor.

10

A solid dispersion substantially as hereinbefore described may also be used in combination with other therapeutic agents which may be useful in the treatment of the above-mentioned disease states. The invention thus provides, in another aspect, a combination of a solid dispersion substantially as hereinbefore described together with another therapeutically active agent, for simultaneous, separate, or sequential use

15

The combination referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical compositions comprising a combination as defined above together with a pharmaceutically acceptable diluent or carrier comprise a further aspect of the invention.

20

The individual components of such a combination may also be administered either sequentially or simultaneously in separate pharmaceutical formulations.

25

Appropriate doses of known therapeutic agents for use in combination with a solid dispersion of the invention will be readily appreciated by those skilled in the art.

30

Compound B may be prepared by any suitable method known in the art, for example as described in WO93.14074.

Compound A may be prepared by any suitable method known in the art, or as substantially hereinafter described in the accompanying Examples.

35

The synthesis of Compound A and of intermediates useful in the preparation thereof, are illustrated by the following, non-limiting Examples.

Intermediate 1

5 (R)-Na^a-(3,4-Methylenedioxyphenylcarbonyl)-tryptophan methyl ester

To a suspension of D-tryptophan methyl ester hydrochloride (10.2g) in anhydrous CH₂Cl₂ (150ml) cooled at 0°C was added dropwise triethylamine (12.3ml). To the resulting solution solid piperonyl chloride (8.16g) was added portionwise at the same temperature, and the mixture was stirred at room temperature for 2 hours. The mixture was washed successively with water, 0.5N hydrochloric acid, water, a saturated aqueous solution of NaHCO₃ and again with water. After drying over Na₂SO₄ and evaporation of the solvent under reduced pressure, the resulting oil on trituration from hot cyclohexane afforded the title compound as a white solid (14.7g).

15 mp : 123-124°C

20°

[α]_D = - 84.4° (c = 1.04, CHCl₃).

20 Intermediate 2

(R)-Na^a-(3,4-Methylenedioxyphenylthiocarbonyl)-tryptophan methyl ester

A mixture of Intermediate 1 (14g) and Lawesson's reagent (9.28g) in dimethoxyethane (280ml) was heated at 60°C under N₂ for 16 hours with stirring. The reaction mixture was evaporated to dryness and the resulting oil was dissolved in ethyl acetate, then washed successively with an aqueous saturated solution of NaHCO₃ and water and dried over Na₂SO₄. The oily residue obtained after evaporation under reduced pressure gave, on trituration from cyclohexane, a yellow powder which was filtered and washed with cooled methanol to afford the title compound (9.74g).

30 mp : 129-130°C

20°

[α]_D = - 186.8° (c = 1.14, CHCl₃).

Intermediate 3

(1R,3R)-Methyl 1,2,3,4-tetrahydro-1-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

A solution of Intermediate 2 (9g) and methyl iodide (10ml) in anhydrous dichloromethane (200ml) was heated at reflux under an argon atmosphere with protection from light. After 24 hours, the solvent was removed under reduced pressure to give an orange oil which on trituration from hexane gave a solid which was washed with ether and used without further purification in the next step. This compound (13.11g) was dissolved in methanol (250ml) and the solution was cooled to -78°C. NaBH₄ (0.99g) was then added by portions and the mixture was stirred at the same temperature for 1 hour. The reaction was quenched by addition of acetone (10ml) and the solvent was removed under reduced pressure. The residue was dissolved in CH₂Cl₂, washed with water and then with brine and dried over Na₂SO₄. After evaporation of the solvent, the orange oil gave on trituration from a hot mixture of diethyl ether/cyclohexane an orange powder which was recrystallised from diethyl ether/pentane to afford the title compound as a pale yellow solid (5.15g).

mp: 154 - 155°C

20°

[α]_D²⁰ = + 24.4° (c = 1.03, CHCl₃).

Intermediate 4

(1R,3R)-Methyl 1,2,3,4-tetrahydro-2-chloroacetyl-(3,4-methylenedioxyphenyl)-9H-pyrido[3,4-b]indole-3-carboxylate

Method A

To a stirred solution of Intermediate 3 (9.7g) and NaHCO₃ (2.79g) in anhydrous CHCl₃ (200ml) was added dropwise chloroacetyl chloride (5.3ml) at 0°C under N₂. The resulting mixture was stirred for 1 hour at the same temperature and diluted with CHCl₃ (100ml). Water (100ml) was then added dropwise with stirring to the mixture, followed by a saturated aqueous solution of NaHCO₃. The organic layer was washed with water until neutrality and dried over Na₂SO₄. After evaporation of the solvent under reduced pressure, the oily compound obtained was crystallised from ether to give the title compound as a pale yellow solid (9.95g).

mp : 233°C

20°
[a]_D = - 125.4° (c = 1.17, CHCl₃).

Method B

- 5 Chloroacetyl chloride (4ml) was added dropwise to a solution of Intermediate 3 (16.1g) and triethylamine (7ml) in anhydrous CH₂Cl₂ (200ml) at 0°C under N₂. The solution was stirred at 0°C for 30 minutes, then diluted with CH₂Cl₂ (300ml). The solution was washed with water (200ml), a saturated aqueous solution of NaHCO₃ (300ml) and brine (400ml). After drying over Na₂SO₄ and
10 evaporation under reduced pressure, the resulting solid was washed with ether (300ml) to give the title compound as a pale yellow solid (18.3g).

Example 1

- 15 (6R,12aR)-2,3,6,7,12,12a-Hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione (Compound A)
To a stirred suspension of Intermediate 4 (12.5g) in MeOH (400ml) was added at room temperature a solution of methylamine (33% in EtOH) (13.7ml) and the resulting mixture was heated at 50°C under N₂ for 14 hours. The solvent was removed under reduced pressure and the residue was dissolved in CH₂Cl₂ (1l).
20 After washing with water (3 x 500ml), drying over Na₂SO₄ and evaporating to dryness, the white solid obtained was recrystallised from 2-propanol to give the title compound as white needles (7.5g).
mp : 298-300°C.

- 25 20°
[a]_D = + 71.3° (c = 0.55, CHCl₃).
Elemental analysis (C₂₂H₁₉N₃O₄) calculated: C, 67.86; H, 4.92; N, 10.79;
found: C, 67.79; H, 4.95; N, 10.61%.

- 30 Co-precipitation of Compound A :Hydroxypropyl methylcellulose phthalate using acetone/water

	%
Compound A	25 - 90
Hydroxypropyl methylcellulose phthalate (HPMCP)*	10 - 75
35 * Grades HP55 and HP50.	

15

Compound A (1g) and HPMCP (1g) were dissolved in a 9:1 mixture of acetone/water (27 ml). 0.25M Hydrochloric acid (83ml) was added. The resultant co-precipitate was filtered, washed with water (5x 3ml), dried in vacuo and milled.

5

Co-precipitation of Compound A :Hydroxypropyl methylcellulose phthalate using tetrahydrofuran/water

	%
Compound A	25 - 90
10 Hydroxypropyl methylcellulose phthalate (HPMCP)*	10 - 75
* Grades HP55 and HP50.	

Compound A (1g) and HPMCP (1g) were dissolved in a 9:1 mixture of tetrahydrofuran/water (10ml) and the resultant solution was added to a solution of glacial acetic acid (2.25ml) in water (37.7ml). The resultant precipitate was filtered, washed with water (5x3ml), dried in vacuo and milled.

15

Co-precipitation of Compound B :Hydroxypropyl methylcellulose phthalate using acetone/methanol

	%
Compound B	25 - 90
20 Hydroxypropyl methylcellulose phthalate (HPMCP)*	10 - 75
* Grades HP55 and HP50.	

Compound B (0.65g) and HPMCP (0.65g) were dissolved in a 1:1 mixture of acetone and methanol (35ml). 0.5N Hydrochloric acid (50ml) was added followed by distilled water (50ml). The resultant precipitate was collected, washed with aqueous hydrochloric acid, filtered, and dried in vacuo.

25

TABLETS FOR ORAL ADMINISTRATION

Co-precipitates of Compound A: HPMCP and Compound B: HPMCP were formulated as follows:

30

Compound A: HPMCP co-precipitate was blended with the excipients. The resultant mix was compressed into tablets.

1.	mg/tablet
Compound A: HPMCP co-precipitate	100.0
Microcrystalline cellulose	289.2
Colloidal silicon dioxide	0.8
Crospovidone	8.0
Magnesium stearate	2.0

2.	mg/tablet
Compound A: HPMCP co-precipitate	100.0
Microcrystalline cellulose	229.2
Lactose (anhydrous)	52.0
Colloidal silicon dioxide	0.8
Crospovidone	16.0
Magnesium stearate	2.0

5

3.	mg/tablet
Compound A: HPMCP co-precipitate	100.0
Microcrystalline cellulose	249.2
Polyvinyl pyrrolidone	40.0
Colloidal silicon dioxide	0.8
Crospovidone	8.0
Magnesium stearate	2.0

4.	mg/tablet
Compound A: HPMCP co-precipitate	100.0
Microcrystalline cellulose	281.2
Sodium lauryl sulphate	8.0
Colloidal silicon dioxide	0.8
Crospovidone	8.0
Magnesium stearate	2.0

17

5.	mg/tablet
Compound A: HPMCP co-precipitate	100.0
Microcrystalline cellulose	61.87
Dibasic calcium phosphate anhydrous	62.00
Croscarmellose Sodium	10.00
Sodium lauryl sulphate	5.0
Polyvidone 30	9.38
Colloidal silicon dioxide	0.5
Magnesium stearate	1.25

Compound B: HPMCP co-precipitate was blended with the excipients. The resultant mix was compressed into tablets.

5

1.	mg/tablet
Compound B: HPMCP co-precipitate	10.0
Microcrystalline cellulose	66.74
Dibasic calcium phosphate anhydrous	66.74
Croscarmellose sodium	6.00
Magnesium stearate	0.53

Tablets of other strengths may be prepared by altering the ratio of Compound A or B: HPMCP co-precipitate to the other excipients.

10

FILM COATED TABLETS

The aforementioned tablet formulations were film coated.

Coating Suspension	% w/w
Opadry white +	13.2
Purified water	to 100.0*

15

* The water did not appear in the final product. The maximum theoretical weight of solids applied during coating was 20mg/tablet.

+ Opadry white is a proprietary material obtainable from Colorcon Limited, U.K. which contains hydroxypropyl methylcellulose, titanium dioxide and triacetin.

- 5 The tablets were film coated using the coating suspension in conventional film coating equipment.

CAPSULES

1.	mg/capsule
Compound A: HPMCP co-precipitate	100.0
Lactose	168.5
Polyvinyl pyrrolidone	30.0
Magnesium stearate	1.5

- 10 The Compound A: HPMCP co-precipitate was blended with the excipients. The mix was filled into size No. O hard gelatin capsules using suitable equipment.

2.	mg/capsule
Compound A: HPMCP co-precipitate	100.0
Microcrystalline cellulose	183.5
Sodium lauryl sulphate	6.0
Crospovidone	9.0
Magnesium stearate	1.5

- 15 The Compound A: HPMCP co-precipitate was blended with the excipients. The mix was filled into size No. O hard gelatin capsules using suitable equipment.

3.	mg/capsule
Compound B: HPMCP co-precipitate	10.0
Lactose	90.0
Microcrystalline cellulose	90.0
Crospovidone	7.0
Sodium lauryl sulphate	1.0

Magnesium stearate	2.0
--------------------	-----

- 5 The compound B:HPMCP co-precipitate was blended with the excipients. The mix was filled into size No. 1 hard gelatin capsules using suitable equipment. Other strengths may be prepared by altering the ratio of Compound A or B to excipient, the fill weight and if necessary changing the capsule size.

CLAIMS

1. A process of preparing a solid dispersion comprising a poorly water soluble drug or salts or solvates thereof, and a pharmaceutically acceptable carrier or excipient therefor, which process comprises:

(a) providing an intimate mixture comprising the carrier or excipient and a non-aqueous, water miscible solvent or combination of solvents, and optionally, water;

(b) co-mixing the intimate mixture obtained in step (a) with a poorly water soluble drug; and

(c) co-precipitating the poorly water soluble drug and the carrier or excipient.

2. A process as claimed in Claim 1, wherein the carrier or excipient is a polymeric material selected from hydroxypropyl methyl cellulose phthalate, polymethylacrylate, and hydroxypropyl cellulose.

3. A process as claimed in Claim 1 or Claim 2, wherein the co-precipitation medium is selected from a weakly acidic, neutral, or basic medium.

4. A process as claimed in any one of Claims 1-3, wherein the non-aqueous, water miscible solvent employed is selected from acetone, methanol, dimethylacetamide, dimethylsulphoxide, dimethylformamide, tetrahydrofuran, and combinations thereof.

5. A process as claimed in any one of Claims 1-4, wherein the poorly water soluble drug is (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione.

6. A process as claimed in any one of Claims 1-4, wherein the poorly water soluble drug is (+)-N-[1-(adamantanmethyl)-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro - 1H-1,5benzodiazepin-3-yl]-N'-phenylurea.

7. A process as claimed in Claim 5 wherein the solvent employed is selected from acetone and tetrahydrofuran in a 10% aqueous mixture.
- 5 8. A process as claimed in Claim 6 wherein the solvent employed is a combination of acetone and methanol.
9. A process as claimed in any one of Claims 1-8, wherein the carrier or excipient is hydroxypropyl methyl cellulose phthalate.
- 10 10. A process as claimed in claim 9, wherein the co-precipitation medium is a weakly acidic medium selected from hydrochloric acid and acetic acid.
11. A process as claimed in any one of Claims 1-8, wherein the carrier or
15 excipient is hydroxypropylcellulose.
12. A process as claimed in Claim 11 wherein the co-precipitation medium is water at an elevated temperature such as 70-80°C.
- 20 13. A process as claimed in any one of Claims 1-8, wherein the carrier or excipient is polymethyl acrylate.
14. A process as claimed in Claim 13 wherein the co-precipitation medium is water or dilute alkali.
- 25 15. A process of preparing a solid dispersion comprising (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4- dione or salts or solvates thereof, and a pharmaceutically acceptable carrier or excipient therefor, which process
30 comprises:
- (a) providing an intimate mixture of a poorly water soluble drug selected from Compound A or Compound B, the carrier or excipient therefor and a non-aqueous, water miscible solvent or combination of solvents, and optionally,
35 water; and

(b) co-precipitating the compound and the carrier or excipient.

5 16. A process as claimed in Claim 15 wherein the carrier or excipient is hydroxypropyl methyl cellulose phthalate.

10 17. A process as claimed in Claim 15 or Claim 16, wherein the co-precipitation medium is a weakly acidic medium selected from hydrochloric acid and acetic acid.

18. A process as claimed in any one of Claims 15-17, wherein the solvent employed is selected from acetone and tetrahydrofuran, in a mixture comprising a 10% ratio of water.

15 19. A process of preparing a solid dispersion comprising (+)-N-[1-(adamantanmethyl)-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro - 1H-1,5benzodiazepin-3-yl]-N'-phenylurea or salts or solvates thereof, and a pharmaceutically acceptable carrier or excipient therefor, which process comprises:

20 (a) providing an intimate mixture of a poorly water soluble drug selected from Compound A or Compound B, the carrier or excipient therefor and a non-aqueous, water miscible solvent or combination of solvents, and optionally, water; and

25 (b) co-precipitating the compound and the carrier or excipient.

30 20. A process as claimed in Claim 19 wherein the carrier or excipient is hydroxypropyl methyl cellulose phthalate.

21. A process as claimed in Claim 19 or Claim 20, wherein the acidic medium is 0.5N hydrochloric acid.

35 22. A process as claimed in any one of Claims 19-21, wherein the solvent employed is a combination of acetone and methanol.

23. A solid dispersion consisting of (6R,12aR)-2,3,6,7,12,12a-hexahydro-2-methyl-6-(3,4-methylenedioxyphenyl)-pyrazino[2',1':6,1]pyrido[3,4-b]indole-1,4-dione and a pharmaceutically acceptable carrier or excipient therefor.

5

24. A solid dispersion as claimed in Claim 23 for use in therapy.

25. A solid dispersion as claimed in Claim 23 for use in the manufacture of a medicament for the treatment of conditions where inhibition of PDEV is of therapeutic benefit.

10

26. A method of treating in conditions where inhibition of PDEV is of therapeutic benefit in a human or non-human animal body which comprises administering to said body a therapeutically effective amount of a solid dispersion as claimed in Claim 23.

15

27. A solid dispersion consisting of (+)-N-[1-(adamantanmethyl)-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro - 1H-1,5benzodiazepin-3-yl]-N'-phenylurea and a pharmaceutically acceptable carrier or excipient therefor.

20

28. A solid dispersion as claimed in Claim 27 for use in therapy.

29. The use of a solid dispersion as claimed in Claim 27 for the manufacture of a medicament for the treatment of conditions where modification of the effects of gastrin and/or CCK-B is of therapeutic benefit.

25

30. A method for the treatment of a mammal, including man, in particular in the treatment of conditions where modification of the effects of gastrin and/or CCK-B is of therapeutic benefit which method comprises administering to the patient an effective amount of a solid dispersion as claimed in Claim 27.

30

31. A pharmaceutical composition comprising a solid dispersion as claimed in Claim 23 or Claim 24, together with a pharmaceutically acceptable carrier therefor.

35

32. A process of preparing a pharmaceutical composition comprising a solid dispersion as claimed in Claim 23 or Claim 24, which process comprises mixing said solid dispersion with a pharmaceutically acceptable carrier therefor.
- 5 33. A combination of a solid dispersion as claimed in Claim 23 or Claim 24, together with another therapeutically active agent, for simultaneous, separate, or sequential use.
- 10 34. A pharmaceutical composition comprising a combination as claimed in Claim 33 together with a pharmaceutically acceptable diluent or carrier.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 96/02299

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K9/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 297 866 (THE BOOTS COMPANY PLC) 4 January 1989 see the whole document ---	1,2,4,11
X,P	WO,A,95 19978 (LABORATOIRES GLAXO) 27 July 1995 see page 32; example 1 see page 71, line 25 - page 74, line 16 ---	23-26, 31,32
X	EP,A,0 558 104 (GLAXO SPA) 1 September 1993 cited in the application see page 40; example 45 see page 41, line 40 - page 43, line 30 -----	27-30

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

30 September 1996

Date of mailing of the international search report

18.10.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patendaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Ventura Amat, A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 96/02299

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-297866	04-01-89	NONE	
WO-A-9519978	27-07-95	AU-A- 1574895	08-08-95
		FI-A- 962927	19-07-96
		ZA-A- 9500424	27-09-95
EP-A-558104	01-09-93	AP-A- 369	06-11-94
		AU-B- 661749	03-08-95
		AU-A- 3193593	22-07-93
		AU-A- 3450193	03-08-93
		BG-A- 98881	31-03-95
		CA-A- 2087672	22-07-93
		CN-A- 1074678	28-07-93
		CZ-A- 9401736	13-09-95
		WO-A- 9314074	22-07-93
		FI-A- 943421	19-07-94
		HU-A- 67375	28-03-95
		JP-A- 7118244	09-05-95
		NO-A- 942720	20-07-94
		NZ-A- 245715	26-07-95
		OA-A- 9964	11-12-95
		SK-A- 86494	08-03-95
		ZA-A- 9300375	18-07-94